

O2k manual titrations: SUIT protocols with mitochondrial preparations

Oroboros Instruments, High-Resolution FluoRespirometry
Schöpfstr 18, A-6020 Innsbruck, Austria
instruments@oroboros.at; www.oroboros.at



O2k chamber volume: 2.0 ml

| Substrates | Event | Concentration in syringe (solvent) | Storage [°C] | Final conc. in 2 ml | Titration [μ l] | Syringe [μ l] |
|-------------------------------------|-------|------------------------------------|--------------|-----------------------------|----------------------|--------------------|
| Pyruvate | P | 2 M (H_2O) | fresh | 5 mM | 5 | 25 |
| Malate | M | 0.4 M (H_2O) | -20 | 2 mM | 10 | 25 |
| Glutamate | G | 2 M (H_2O) | -20 | 10 mM | 10 | 25 |
| Succinate* | S | 1 M (H_2O) | -20 | 10 mM | 20 | 50 |
| Octanoyl carnitine | Oct | 0.1 M (H_2O) | -20 | 0.2 mM | 4 | 10 |
| Ascorbate | As | 0.8 M (H_2O) | -20 | 2 mM | 5 | 25 |
| TMPD | Tm | 0.2 M (H_2O) | -20 | 0.5 mM | 5 | 25 |
| Cyt. c | c | 4 mM (H_2O) | -20 | 10 μ M | 5 | 25 |
| ADP+ Mg ²⁺ | D | 0.5 M (H_2O) | -80 | 1 - 5 mM | 4 - 20 | 25 |
| ATP+ Mg ²⁺ | T | 0.5 M (H_2O) | -80 | 1 - 5 mM | 4 - 20 | 25 |
| Uncoupler | | | | | | |
| CCCP‡ | U | 0.1 mM (EtOH) | -20 | 0.05 μ M steps | 1 μ l steps | 10 |
| CCCP‡ | U | 1.0 mM (EtOH) | -20 | 0.5 μ M steps | 1 μ l steps | 10 |
| Inhibitors | | | | | | |
| Rotenone | Rot | 1 mM (EtOH) | -20 | 0.5 μ M | 1 | 10 |
| Malonic acid | Mna | 2 M (H_2O) | fresh | 5 mM | 5 | 25 |
| Antimycin A | Ama | 5 mM (EtOH) | -20 | 2.5 μ M | 1 | 10 |
| Myxothiazol | Myx | 1 mM (EtOH) | -20 | 0.5 μ M | 1 | 10 |
| Sodium azide | Azd | 4 M (H_2O) | -20 | \geq 100 mM | \geq 50 | 50 |
| KCN | Kcn | 1 M (H_2O) | fresh | 1.0 mM | 2 | 10 |
| Oligomycin | Omy | 5 mM (EtOH) | -20 | 2.5 μ M | 1 | 10 |
| Carboxyatractyloside | Cat | 5 mM (H_2O) | -20 | 5 μ M | 2 | 10 |
| Other | | | | | | |
| Digitonin | Dig | 10 mg/ml (DMSO) | -20 | 10μ g· 10^{-6} cells | 1 μ l 10^{-6} | 10 |
| Catalase in MiR06 | Ctl | 112,000 U/ml | -20 | 280 U/ml | 5 | 25 |
| Hydrogen peroxide for reoxygenation | Hp | 200 mM | fresh | | 1 - 3 | 10 |

* The concentration of S may be increased up to 50 mM after Rot to compensate for the inhibitory effect of M.

‡ 0.1 mM stock for mt-preparations with high uncoupler sensitivity; 1 mM stock for mt-preparations with low uncoupler sensitivity, intact cells in various culture media (e.g. RPMI, DMEM, EGC) and for TIP2k.

O2k-Chamber volume: 2.0 ml

| Fluorescence probes and related | Event | Concentration in syringe (solvent) | Storage [°C] | Final conc. in 2 ml | Titration [μ l] | Syringe [μ l] |
|-----------------------------------|-------|------------------------------------|--------------|---------------------|----------------------|--------------------|
| DTPA | DTPA | 5 mM (H ₂ O) | -20 | 15 μ M | 6 | 20 |
| Amplex®UltraRed | AmR | 10 mM (DMSO) | -20 | 10 μ M | 2 | 10 |
| Horse radish peroxidase | HRP | 500 U/ml (MiR05) | -20 | 1 U/ml | 4 | 10 |
| Superoxide dismutase | SOD | check supplier information | 4-8 | 5 U/ml | | 10 |
| Hydrogen peroxide for calibration | Hp | 0.04 mM (H ₂ O) | fresh | 0.1 μ M | 5 | 10 |
| Safranin | Saf | 0.2 mM (H ₂ O) | RT | 0.25 μ M | 2.5 | 10 |
| TMRM | Tmr | 0.2 mM (H ₂ O) | -20 | 0.25 μ M | 2.5 | 10 |
| Calcium Green | CaG | 2 mM (H ₂ O) | -20 | 1 μ M | 1 | 10 |
| Magnesium Green | MG | 5 mM (H ₂ O) | -20 | 2.5 μ M | 1 | 10 |

Further abbreviations

| | |
|--|------------------|
| Attractyloside | Atr |
| Calcium | Ca ²⁺ |
| Dinitrophenol | DNP; U |
| Diethylenetriamin- | |
| N,N,N',N,N- | |
| pentaacetic acid | DTPA |
| Carbonyl cyanide p-trifluoro-methoxyphenyl hydrazone | FCCP; U |
| Glucose | Glc |
| Glycerophosphate | Gp |
| Hydroxycinnamate | Hci |
| Oxaacetate | Oa |
| Octanoate | Oca; FA |
| Palmitate | Paa; FA |
| Palmitoylcarnitine | Pal; FA |
| Tetraphenyl-phosphonium ion | TPP ⁺ |

References

- Chinopoulos C, Kiss G, Kawamata H, Starkov AA (2014) Measurement of ADP-ATP exchange in relation to mitochondrial transmembrane potential and oxygen consumption. Methods Enzymol 542:333-48. »[Bioblast link](#)«
- Elustondo PA, Negoda A, Kane CL, Kane DA, Pavlov EV (2014) Spermine selectively inhibits high-conductance, but not low-conductance calcium-induced permeability transition pore. Biochim Biophys Acta 1847:231-40. »[Bioblast link](#)«
- Gnaiger E (2014) Mitochondrial pathways and respiratory control. An introduction to OXPHOS analysis. 4th ed. Mitochondr Physiol Network 19.12. Oroboros MiPNet Publications, Innsbruck:80 pp. »[Bioblast link](#)«
- Krumschnabel G, Eigenthaler A, Fasching M, Gnaiger E (2014) Use of safranin for the assessment of mitochondrial membrane potential by high-resolution respirometry and fluorometry. Methods Enzymol 542:163-81. »[Bioblast link](#)«
- Krumschnabel G, Fontana-Ayoub M, Sumbalova Z, Heidler J, Gauper K, Fasching M, Gnaiger E (2015) Simultaneous high-resolution measurement of mitochondrial respiration and hydrogen peroxide production. Methods Mol Biol 1264:245-61. »[Bioblast link](#)«
- Pesta D, Gnaiger E (2012) High-resolution respirometry. OXPHOS protocols for human cells and permeabilized fibres from small biopsies of human muscle. Methods Mol Biol 810: 25-58. »[Bioblast link](#)«